

Frequency:

Continuous up to 24 hours

Starting

7/13/2023



Today

Tomorrow

For

24



Hours

Days

Weeks

At

1150



Starting: **Today 1150** Ending: **Tomorrow**

- ! Reason for restraint: Support patient safety Imminent risk of patient physically harming self or others Rational for use
- ! The behavior exhibited that required the application of Restraints/Seclusion:

 - Hitting, kicking, throwing objects, biting, etc...
 - Self-harmful behavior (head banging, thrashing, self mutilation)
 - Violent/Aggressive/Assaultive Threatens imminent harm to self or others Other
- ! Alternative evaluated or attempted:

 - Verbal de-escalation Opportunity to express feelings Reviewed medications Assess patient's comfort
 - Patient allowed to ambulate when possible Physical needs met Patient reoriented, comforted, reassured
 - Diversionary activities attempt environment alteration Family involvement
 - Moved the patient closer to the nursing station Rounding for safety Other
- Type of restraint: Mitten Limb restraints Vest **Net Bed** Seclusion (VPH only) Physical hold (Typically used in VPH)
- ! Discontinuation goal: No hitting, kicking, throwing objects, biting, etc...

 - No self harmful behavior (head banging, thrashing, self mutilation) No violent/aggressive/assaultive behavior
 - No threats of imminent harm to self or other Other

Violent restraint assessment & provider discussion required at specified timeframes:

Child less than 9 years old = Q1HR

Child 9 - 17 years old = Q2HR

Patient greater than or equal to 18 years old = Q4HR

Comments:

+ [Add Comments](#)

Reference Links:

- [Restraint Order Discontinuation Protocol](#)

✓ Accept

✗ Cancel

Link Order

— Remove

Restraints Non-Violent or Non-Self Destructive Panel

✓ Accept

Frequency:

Continuous up to 48 hours

Starting

7/13/2023



Today

Tomorrow

For

48



Hours

Days

Weeks

At

1150



Starting: **Today 1150** Ending: **Sat 7/15**

Reason for restraint: Support patient safety Provide optimum medical care To protect medically necessary device(s)

Behavior indicating need for restraint:

Agitation Confused / Disoriented

Uncooperative with necessary medical care (attempts to pull out i.e. ETT, feeding tubes, intravenous lines)

Unable to follow directions Other

Alternative evaluated or attempted:

Verbal de-escalation Opportunity to express feelings Reviewed medications Assess patient's comfort

Patient allowed to ambulate when possible Physical needs met Patient reoriented, comforted, reassured

Diversionary activities attempt environment alteration Family involvement

Moved the patient closer to the nursing station Rounding for safety Other

Type of restraint:

Mitten Limb restraints Vest Net Bed Roll Belt

Discontinuation goal: Patient is no longer agitated Patient is no longer confused / disoriented

Patient is no longer uncooperative with necessary medical care (attempts to pull out i.e. ETT, feeding tubes, intravenous lines)

Patient is able to follow directions Other